

Hoar SK, et al., Agricultural herbicide use and risk of lymphoma and soft-tissue sarcoma. JAMA 256:1141-7; 1986

Abstract:

A population-based case-control study of soft-tissue sarcoma (STS), Hodgkin's disease (HD), and non-Hodgkin's lymphoma (NHL) in Kansas found farm herbicide use to be associated with NHL (odds ratio [OR], 1.6; 95% confidence interval [CI], 0.9, 2.6). Relative risk of NHL increased significantly with number of days of herbicide exposure per year and latency. Men exposed to herbicides more than 20 days per year had a six fold increased risk of NHL (OR, 6.0; 95% CI, 1.9, 19.5) relative to non-farmers. Frequent users who mixed or applied the herbicides themselves had an OR of 8.0 (95% CI, 2.3, 27.9) for NHL. Excesses were associated with use of phenoxyacetic acid herbicides, specifically 2,4-dichlorophenoxyacetic acid. Neither STS nor HD was associated with pesticide exposure. This study confirms the reports from Sweden and several US states that NHL is associated with farm herbicide use, especially phenoxyacetic acids. It does not confirm the case-control studies or the cohort studies of pesticide manufacturers and Vietnam veterans linking herbicides to STS or HD.

Erratum in: JAMA 1986 Dec 26;256(24):3351